

# Transgenerational inheritance of DNA hypomethylation in *Daphnia magna* in response to salinity stress

Guilherme Jeremias<sup>1</sup> • João Barbosa<sup>2</sup> • Sérgio M. Marques<sup>1</sup> • Karel De Schamphelaere<sup>2</sup> • Filip Van Nieuwerburgh<sup>3</sup> • Dieter Deforce<sup>3</sup> • Fernando J. M. Gonçalves<sup>1</sup> • Joana Luísa Pereira<sup>1</sup> ✉ jpereira@ua.pt • Jana Asselman<sup>2</sup>

<sup>1</sup> Department of Biology & CESAM, University of Aveiro, Portugal • <sup>2</sup> Laboratory for Environmental Toxicology and Aquatic Ecology (GhEnToxLab), Ghent University, Belgium • <sup>3</sup> Laboratory of Pharmaceutical Biotechnology, Ghent University, Belgium

## Introduction


- Epigenetic mechanisms play an important role in environmental stress response and regulation.
- Epigenetic marks can be inherited through successive generations, yet few studies have shown persisting stress-induced transgenerational effects.
- Daphnia* is a parthenogenetic model and key species in freshwater ecosystems, widely considered among regulatory frameworks.

## Aim

To explore the potential for transgenerational inheritance (maintained up to unexposed F3) of DNA methylation patterns in *Daphnia* after F0 exposure to salinity and the underlying molecular mechanisms.


## Material & Methods

In addition to F0, the somatic and/or germline cells of developing embryos can also be directly exposed to the parental stressor



➔ So... DNA methylation patterns were monitored until F3 generation to spot “true” transgenerational effects

Six monoclonal cultures of 70 neonates (0 g/L NaCl (x 3) and 4.1 g/L NaCl (x 3) in ASTM medium)



+ and - indicate the history of each culture regarding exposure to NaCl (+) and maintenance in clean medium (-)

Red boxes: treatments from which DNA was extracted

### DNA extraction

MasterPure Complete DNA and RNA Purification Kit (Epicenter, Madison, WI, USA)

Reduced Representation Bisulfite Sequencing

Bioinformatic and Statistical analysis

FastQC; Trim Galore!; Bismark; R software


### Life history traits

age at first reproduction, net reproductive rate, per capita rate of population increase


## Results

7 genes significantly differentially methylated across the four generations (F0–F1–F2–F3)

Venn Diagram overlapping the genes differentially methylated between NaCl treatments and the control through the different generations (p<0.05)



Differences in methylation (in black) between the control and F0-F3 salinity treatment. Control treatment methylation is plotted in orange



No main effects of NaCl across all generations for the per capita rate of population increase, age at first reproduction or net reproductive rate

## Concluding remarks

- The exposure of a single *D. magna* generation to high levels of salinity (F0) provoked specific methylation patterns that were transferred to the three non-exposed generations (F1, F2, F3).
- Epigenetic changes seem to be particularly targeted to genes involved in coping with general cellular stress responses.
- No life history costs seemed to be associated with the maintenance of hypomethylation across the unexposed generations.

Epigenetic marks can be useful indicators of past or historic pollution in *Daphnia*

### Literature cited

Brander SM, Biales AD, Connon RE (2017) The Role of Epigenomics in Aquatic Toxicology. Environ Toxicol Chem. 36:2565–2573. doi: 10.1002/etc.393



### Acknowledgements

Thanks are due for the financial support to CESAM (UID/AMB/50017/2019), to FCT/MCTES through national funds. G Jeremias is recipient of an individual research grants by FCT (ref SFRH/BD/139076/2018). J.L Pereira and S Marques are funded by national funds (OE), through FCT, in the scope of the framework contract foreseen in article 23 of the Decree-Law 57/2016 changed by Law 57/2017. J.A. is a postdoctoral fellow of the Science Foundation Flanders (FWO Vlaanderen) and was the recipient of an individual research grant by FWO (1521617N).